

page 10, second last full paragraph to page 11, page 21, first full paragraph, page 27, second full paragraph, and originally filed claim 20. Additionally, these independent claims are related to claims 1-15 of Issued Patent No. 5,932,540.

Moreover, support for representative claims 34-35, can be found, for example, at page 18, first full paragraph and page 27, first full paragraph. Support for representative claims 36-39 can be found, for example, at page 23, second full paragraph and page 26, fourth full paragraph. Support for representative claims 40-41 can be found, for example, at page 26, fifth full paragraph. Additionally, support for representative claim 42 can be found, for example, at page 11, last full paragraph that extends to page 12. Support for representative claim 44 can be found, for example, at page 32, last full paragraph to page 33.

Furthermore, support for representative claims 44-53 can be found, for example, at pages 27, last full paragraph, to page 28, and are related to claims 61-185 of Issued Patent No. 5,932,540.

Thus, no new matter has been added by way of the amendment.

I. Amendment of the Specification.

The specification has been amended to correct an obvious typographical error. 5x SSC is a well-known solution used in hybridization solutions. (*See, e.g.,* Exhibit A, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, N.Y., at page 2.10.7 (1989).) SSC is normally made as a 20x stock solution, and then diluted accordingly for a particular use. Exhibit B shows that a 20x SSC stock solution contains 3 M NaCl and 0.3 M trisodium citrate. (*See, e.g.,* Exhibit B, CURRENT PROTOCOLS, at page A.2.5.) To make a 5x SSC solution, the 20x solution must be diluted by a factor of four. Therefore, a 5x SSC solution contains 750 mM NaCl ($3\text{ M} \div 4 = 750\text{ mM}$) and 75 mM trisodium citrate ($0.3\text{ M} \div 4 = 75\text{ mM}$). One skilled in the art would have immediately recognized that the amount of ingredients listed in the specification for a 5x SSC solution was incorrect. Rather than describing a 5x SSC solution, made up of 750 mM NaCl and 75 mM trisodium citrate, the specification inaccurately listed the ingredient amounts for a 1x solution. The skilled artisan, in recognizing the typographical error, could have easily adjusted the amount of ingredients described in the specification to properly make a 5x SSC solution.

Therefore, because no new matter will be added to the specification if these typographical errors are corrected, Applicants respectfully request that the amendments to the specification to recite the correct ingredient amounts in 5x SSC be entered.

II. ATCC Deposit

To demonstrate full compliance with 37 C.F.R. §§ 1.803-1.809 and to satisfy the requirement of 35 U.S.C. § 112, first paragraph, Applicants assure the Examiner that ATCC Deposit No. 97149 has been deposited under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure with the following International Depository Authority: American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, USA. The deposit comprise a recombinant nucleic acid vector into which cDNA sequence encoding Vascular Endothelial Growth Factor 2 (VEGF-2) has been inserted. The deposit was made on May 12, 1995 as disclosed on page 8, lines 1-3 of the instant application.

In accordance with MPEP § 2410.01 and 37 C.F.R. § 1.808, assurance is hereby given that all restrictions on the availability to the public of the above nucleic acid molecule encoding human VEGF-2, will be irrevocably removed upon the grant of a patent based on the captioned application, and that the deposit will be replaced if viable samples cannot be dispensed by the ATCC, except as permitted under 37 C.F.R. § 1.808(b).

III. The Restriction Requirement.

The Examiner has required an election under 35 U.S.C. § 121 of one of Groups I-XII. In response, Applicants provisionally elect, *with traverse*, Group II represented by claims new claims 33-346 for further prosecution. Applicants reserve the right to file one or more divisional applications directed to non-elected inventions should the restriction requirement be made final.

Applicants respectfully request that the Examiner enter the following amendments prior to examination of the captioned application.

Applicants respectfully traverse the restriction requirement as it applies to Groups I and III-XII. As the Examiner points out, polynucleotides, polypeptides, antibodies, etc. are

patentably distinct inventions. However, even where two patentably distinct inventions appear in a single application, restriction remains improper unless it can be shown that the search and examination of both groups would entail a "serious burden". *See*, M.P.E.P. § 803.

In the present situation, no such showing has been made. Indeed, no arguments have been made explaining why it would impose an undue burden to examine Groups I-XII together.

Applicants submit that a search of the polypeptide claims would provide useful information for Groups I and III-XII. For example, in many if not most publications, where a published polypeptide sequence, the authors also routinely include a description of the polynucleotides and antibodies. Thus, the searches for polypeptides, polynucleotides, and antibodies, etc. commonly overlap. Thus, the search and examination of a polypeptide sequence, and remaining groups would not entail a serious burden. Thus, the searches for Groups I-XII would be overlapping.

Accordingly, as applied to Groups I-XII, the restriction requirement should be withdrawn.


Conclusion

In view of the foregoing remarks, applicants believe that this application is now in condition for allowance.

If there are any fees due in connection with the filing of this paper, please charge the fees to our Deposit Account No. 08-3425. If a fee is required for an extension of time under 37 C.F.R. § 1.136 not accounted for above, such an extension is requested and the fee should also be charged to our Deposit Account.

Respectfully submitted,

Dated: 12/15/99


Michele M. Wales (Reg. No. 43,975)
Attorney for Applicants

Human Genome Sciences, Inc.

9410 Key West Avenue
Rockville, MD 20850
Telephone: (301) 610-5772

Enclosures
MMW/lcc

CURRENT PROTOCOLS IN MOLECULAR BIOLOGY

VOLUME 1

EDITORIAL BOARD

Frederick M. Ausubel
Massachusetts General Hospital & Harvard Medical School

Roger Brent
Massachusetts General Hospital & Harvard Medical School

Robert E. Kingston
Massachusetts General Hospital & Harvard Medical School

David D. Moore
Massachusetts General Hospital & Harvard Medical School

J.G. Seidman
Harvard Medical School

John A. Smith
University of Alabama

Kevin Struhl
Harvard Medical School

GUEST EDITORS

Lisa M. Albright
DNA Sequencing

Donald M. Coen
Harvard Medical School
Polymerase Chain Reaction

Ajit Varki
University of California San Diego
Glycoproteins

SERIES EDITOR

Virginia Benson Chanda



John Wiley & Sons, Inc.

CORE 13 (S36)

Copyright © 1994–1997 by John Wiley & Sons, Inc.

Copyright © 1987–1994 by Current Protocols

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

While the authors, editors, and publisher believe that the specification and usage of reagents, equipment, and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of experimental reagents, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each chemical, piece of equipment, reagent, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. This is particularly important in regard to new or infrequently employed chemicals or experimental reagents.

Library of Congress Cataloging in Publication Data:

Current protocols in molecular biology. 3 vols.

1. Molecular biology—Technique. 2. Molecular biology—Laboratory manuals. I. Ausubel, Frederick M.

QH506.C87 1987 574.8'8'028 87-21033

ISBN 0-471-50338-X

Printed in the United States of America

20 19 18 17 16 15 14 13

IN SITU HYBRIDIZATION AND IMMUNOHISTOCHEMISTRY

14

INTRODUCTION	14.0.3
14.1 Fixation, Embedding, and Sectioning of Tissues, Embryos, and Single Cells	14.1.1
Basic Protocol: Paraformaldehyde Fixation and Paraffin Wax Embedding of Tissues and Embryos	14.1.1
Alternate Protocol: Fixation of Suspended and Cultured Cells	14.1.3
Support Protocol: Perfusion of Adult Mice	14.1.4
Support Protocol: Sectioning Samples in Wax Blocks	14.1.4
Reagents and Solutions	14.1.6
Commentary	14.1.7
14.2 Cryosectioning	14.2.1
Basic Protocol: Specimen Preparation and Sectioning	14.2.1
Support Protocol: Fixation of Cryosections for In situ Hybridization	14.2.5
Support Protocol: Tissue Fixation and Sucrose Infusion	14.2.6
Commentary	14.2.6
14.3 In situ Hybridization to Cellular RNA	14.3.1
Basic Protocol: Hybridization Using Paraffin Sections and Cells	14.3.1
Alternate Protocol: Hybridization Using Cryosections	14.3.5
Support Protocol: Synthesis of ³⁵ S-Labeled Riboprobes	14.3.7
Support Protocol: Synthesis of ³⁵ S-Labeled Double-Stranded DNA Probes	14.3.8
Reagents and Solutions	14.3.8
Commentary	14.3.11
14.4 Detection of Hybridized Probe	14.4.1
Basic Protocol: Film Autoradiography	14.4.1
Basic Protocol: Emulsion Autoradiography	14.4.1
Support Protocol: Preparation of Diluted Emulsion for Autoradiography	14.4.2
Commentary	14.4.3
14.5 Counterstaining and Mounting of Autoradiographed In situ Hybridization Slides	14.5.1
Basic Protocol: Giemsa Staining	14.5.1
Alternate Protocol: Hematoxylin/Eosin Staining	14.5.2
Alternate Protocol: Toluidine Blue Staining	14.5.3
Alternate Protocol: Hoechst Staining	14.5.3
Reagents and Solutions	14.5.4
Commentary	14.5.5
14.6 Immunohistochemistry	14.6.1
Basic Protocol: Immunofluorescent Labeling of Cells Grown as Monolayers	14.6.1
Alternate Protocol: Immunofluorescent Labeling of Suspension Cells	14.6.2
Basic Protocol: Immunofluorescent Labeling of Tissue Sections	14.6.3
Alternate Protocol: Immunofluorescent Labeling of Tissue Sections Using Coplin Jars	14.6.4
Alternate Protocol: Immunofluorescent Labeling Using Streptavidin-Biotin Conjugates	14.6.5
Alternate Protocol: Immunogold Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunoperoxidase Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunofluorescent Double-Labeling of Tissue Sections	14.6.7
Reagents and Solutions	14.6.7
Commentary	14.6.8

continued

- 1c. *Harsh treatment:* Pour several hundred milliliters of boiling 0.1% SDS onto the membrane. Cool to room temperature.

If a membrane is to be reprobbed, it must not be allowed to dry out between hybridization and stripping. If it becomes dry, the probe may bind to the matrix.

2. Place membrane on a sheet of dry Whatman 3MM filter paper and blot excess liquid with a second sheet. Wrap the membrane in plastic wrap and set up an autoradiograph.

If signal is still seen after autoradiography, rewash using harsher conditions.

3. The membrane can now be rehybridized. Alternatively, it can be dried and stored for later use.

Membranes can be stored dry between Whatman 3MM paper for several months at room temperature. For long-term storage, place the membranes in a desiccator at room temperature or 4°C.

REAGENTS AND SOLUTIONS

Aqueous prehybridization/hybridization (APH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see below) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Denatured salmon sperm DNA

Dissolve 10 mg Sigma type III salmon sperm DNA (sodium salt) in 1 ml water. Pass vigorously through a 17-G needle 20 times to shear the DNA. Place in a boiling water bath for 10 min, then chill. Use immediately or store at -20°C in small aliquots. If stored, reheat to 100°C for 5 min and chill on ice immediately before using.

Formamide prehybridization/hybridization (FPH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

50% (w/v) formamide

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see above) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Commercial formamide is usually satisfactory for use. If the liquid has a yellow color, deionize as follows: add 5 g of mixed-bed ion-exchange resin [e.g., Bio-Rad AG 501-X8 or 501-X8(D) resins] per 100 ml formamide, stir at room temperature for 1 hr, and filter through Whatman no. 1 paper.

CAUTION: Formamide is a teratogen. Handle with care.

Labeling buffer

200 mM Tris-Cl, pH 7.5

30 mM MgCl₂

10 mM spermidine

Mild stripping solution

5 mM Tris-Cl, pH 8.0

2 mM EDTA

0.1× Denhardt solution (APPENDIX 2)

CURRENT PROTOCOLS IN MOLECULAR BIOLOGY

VOLUME 1

EDITORIAL BOARD

Frederick M. Ausubel
Massachusetts General Hospital & Harvard Medical School

Roger Brent
Massachusetts General Hospital & Harvard Medical School

Robert E. Kingston
Massachusetts General Hospital & Harvard Medical School

David D. Moore
Massachusetts General Hospital & Harvard Medical School

J.G. Seidman
Harvard Medical School

John A. Smith
University of Alabama

Kevin Struhl
Harvard Medical School

GUEST EDITORS

Lisa M. Albright
DNA Sequencing

Donald M. Coen
Harvard Medical School
Polymerase Chain Reaction

Ajit Varki
University of California San Diego
Glycoproteins

SERIES EDITOR

Virginia Benson Chanda



John Wiley & Sons, Inc.

CORE 13 (S36)

Copyright © 1994–1997 by John Wiley & Sons, Inc.

Copyright © 1987–1994 by Current Protocols

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

While the authors, editors, and publisher believe that the specification and usage of reagents, equipment, and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of experimental reagents, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each chemical, piece of equipment, reagent, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. This is particularly important in regard to new or infrequently employed chemicals or experimental reagents.

Library of Congress Cataloging in Publication Data:

Current protocols in molecular biology. 3 vols.

1. Molecular biology—Technique. 2. Molecular biology—Laboratory manuals. I. Ausubel, Frederick M.

QH506.C87 1987 574.8'028 87-21033
ISBN 0-471-50338-X

Printed in the United States of America

20 19 18 17 16 15 14 13

IN SITU HYBRIDIZATION AND IMMUNOHISTOCHEMISTRY

14

INTRODUCTION	14.0.3
14.1 Fixation, Embedding, and Sectioning of Tissues, Embryos, and Single Cells	14.1.1
Basic Protocol: Paraformaldehyde Fixation and Paraffin Wax Embedding of Tissues and Embryos	14.1.1
Alternate Protocol: Fixation of Suspended and Cultured Cells	14.1.3
Support Protocol: Perfusion of Adult Mice	14.1.4
Support Protocol: Sectioning Samples in Wax Blocks	14.1.4
Reagents and Solutions	14.1.6
Commentary	14.1.7
14.2 Cryosectioning	14.2.1
Basic Protocol: Specimen Preparation and Sectioning	14.2.1
Support Protocol: Fixation of Cryosections for In situ Hybridization	14.2.5
Support Protocol: Tissue Fixation and Sucrose Infusion	14.2.6
Commentary	14.2.6
14.3 In situ Hybridization to Cellular RNA	14.3.1
Basic Protocol: Hybridization Using Paraffin Sections and Cells	14.3.1
Alternate Protocol: Hybridization Using Cryosections	14.3.5
Support Protocol: Synthesis of ³⁵ S-Labeled Riboprobes	14.3.7
Support Protocol: Synthesis of ³⁵ S-Labeled Double-Stranded DNA Probes	14.3.8
Reagents and Solutions	14.3.8
Commentary	14.3.11
14.4 Detection of Hybridized Probe	14.4.1
Basic Protocol: Film Autoradiography	14.4.1
Basic Protocol: Emulsion Autoradiography	14.4.1
Support Protocol: Preparation of Diluted Emulsion for Autoradiography	14.4.2
Commentary	14.4.3
14.5 Counterstaining and Mounting of Autoradiographed In situ Hybridization Slides	14.5.1
Basic Protocol: Giemsa Staining	14.5.1
Alternate Protocol: Hematoxylin/Eosin Staining	14.5.2
Alternate Protocol: Toluidine Blue Staining	14.5.3
Alternate Protocol: Hoechst Staining	14.5.3
Reagents and Solutions	14.5.4
Commentary	14.5.5
14.6 Immunohistochemistry	14.6.1
Basic Protocol: Immunofluorescent Labeling of Cells Grown as Monolayers	14.6.1
Alternate Protocol: Immunofluorescent Labeling of Suspension Cells	14.6.2
Basic Protocol: Immunofluorescent Labeling of Tissue Sections	14.6.3
Alternate Protocol: Immunofluorescent Labeling of Tissue Sections Using Coplin Jars	14.6.4
Alternate Protocol: Immunofluorescent Labeling Using Streptavidin-Biotin Conjugates	14.6.5
Alternate Protocol: Immunogold Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunoperoxidase Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunofluorescent Double-Labeling of Tissue Sections	14.6.7
Reagents and Solutions	14.6.7
Commentary	14.6.8

continued

14.0.1

SDS electrophoresis buffer, 5×

15.1 g Tris base

72.0 g glycine

5.0 g SDS

H₂O to 1000 ml

Dilute to 1× or 2× for working solution, as appropriate

Do not adjust the pH of the stock solution, as the solution is pH 8.3 when diluted. Store at 0° to 4°C until use (up to 1 month).

SED (standard enzyme diluent)

20 mM Tris-Cl, pH 7.5

500 µg/ml bovine serum albumin (Pentax Fraction V)

10 mM 2-mercaptoethanol

Store up to 1 month at 4°C

Sodium acetate, 3 M

Dissolve 408 g sodium acetate·3H₂O in 800 ml H₂O

Add H₂O to 1 liter

Adjust pH to 4.8 or 5.2 (as desired) with 3 M acetic acid

Sodium acetate buffer, 0.1 M

Solution A: 11.55 ml glacial acetic acid/liter (0.2 M).

Solution B: 27.2 g sodium acetate (NaC₂H₃O₂·3H₂O)/liter (0.2 M).

Referring to Table A.2.2 for desired pH, mix the indicated volumes of solutions A and B, then dilute with H₂O to 100 ml. (See Potassium acetate buffer recipe for further details.)

Sodium phosphate buffer, 0.1 M

Solution A: 27.6 g NaH₂PO₄·H₂O per liter (0.2 M).

Solution B: 53.65 g Na₂HPO₄·7H₂O per liter (0.2 M).

Referring to Table A.2.3 for desired pH, mix the indicated volumes of solutions A and B, then dilute with H₂O to 200 ml. (See Potassium phosphate buffer recipe for further details.)

SSC (sodium chloride/sodium citrate), 20×

3 M NaCl (175 g/liter)

0.3 M Na₃citrate·2H₂O (88 g/liter)

Adjust pH to 7.0 with 1 M HCl

STE buffer

10 mM Tris-Cl, pH 7.5

10 mM NaCl

1 mM EDTA, pH 8.0

TAE (Tris/acetate/EDTA) electrophoresis buffer

50× stock solution:

242 g Tris base

57.1 ml glacial acetic acid

37.2 g Na₂EDTA·2H₂O

H₂O to 1 liter

Working solution, pH ~8.5:

40 mM Tris-acetate

2 mM Na₂EDTA·2H₂O

TBE (Tris/borate/EDTA) electrophoresis buffer

10× stock solution, 1 liter:

108 g Tris base (890 mM)

55 g boric acid (890 mM)

40 ml 0.5 M EDTA, pH 8.0 (20 mM)